

Poster #1-40

Warming Increases Plant-Available Nutrients in the SPRUCE Bog

Colleen M. Iversen^{1*}, John Latimer², Deanne J. Brice¹, Joanne Childs¹, Holly M. Vander Stel¹; Natalie A. Griffiths¹, Avni Malhotra¹, Richard J. Norby¹, Keith Olehieser², Steve Sebestyen³, Eric Ward¹, Jeffrey Warren¹, and Paul J. Hanson¹

¹Environmental Sciences Division and Climate Change Science Institute, Oak Ridge National Laboratory, Oak Ridge, TN

²XCEL Engineering, Oak Ridge, TN

³Center for Research on Ecosystem Change, Northern Research Station, USDA Forest Service, Grand Rapids, MN

Contact: iversencm@ornl.gov

BER Program: TES

Project: ORNL Terrestrial Ecosystem Science SFA

Project Website: <https://mnspruce.ornl.gov/>

Peatlands store nearly one-third of global soil carbon in deep deposits of peat that are shielded from decomposition by acidic conditions, waterlogged soils, and cold temperatures. Warming is expected to increase the release of carbon from highly-organic peatland soils, potentially leading to a positive feedback to future warming. This response is expected to be mediated by the response of peatland vegetation to rising atmospheric [CO₂], as well as the effects of warming on plant-available nutrients and water.

We quantified the effects of a range of ecosystem warming (from +0 °C to +9 °C), as well as elevated [CO₂], on plant-available nutrients in the SPRUCE (Spruce and Peatland Responses Under Changing Environments) experiment in an ombrotrophic bog in northern Minnesota, USA. We used ion-exchange resin capsules to monitor monthly changes in plant-available nutrients (i.e., NH₄-N, NO₃-N, and PO₄³⁻) throughout the peat profile and across hummock-hollow microtopography. NH₄-N was by far the most available N source, with NO₃-N making up a negligible fraction; PO₄³⁻ availability was intermediate. After the first full year of increases in temperature and [CO₂], warming—combined with a longer frost-free period—tripled the availability of NH₄⁺ and PO₄³⁻ in the warmest treatment plots. However, the increase in nutrients was much greater below the rooting zone. There is thus far no effect of elevated [CO₂] on nutrient availability.

Interestingly, the same warming response was not apparent in the subset of porewater nutrients collected and measured at bi-weekly intervals at a comparable depth increment in the hollows. While porewater total organic carbon concentrations were increased by warming, indicating increased mineralization of organic matter, there was no difference in porewater NH₄⁺NO₃⁻, or PO₄³⁻ concentrations across the warmed plots.

Taken together, these lines of evidence indicate that warming has increased the mineralization of peat, leading to increased nutrient availability. In turn, increased nutrient uptake by the vegetation has depleted the availability of nutrients in the rooting zone and in porewater. The additional nutrients taken up by the plant community are detectable in increased nitrogen and phosphorus concentrations in Sphagnum mosses as well as vascular plant tissues; however, this may be luxury uptake, as evidence indicates that at least moss production has declined with warming. The relative balance of peat accumulation will depend in part on whether vegetation growth is increased in response to warming and increased nutrient availability, or whether this response is limited by increased drying in the warmed treatments.