

**Poster #9-19****How do Whole-Ecosystem Warming and Elevated Atmospheric Carbon Dioxide Concentrations Affect Peatland Methane Production**

Anya Hopple<sup>1,2</sup>, Rachel Wilson<sup>3</sup>, Glenn Woerdle<sup>1</sup>, Cassandra Zalman<sup>1</sup>, Jeff Chanton<sup>3</sup>, Paul J. Hanson<sup>4</sup>, Jason K. Keller<sup>1\*</sup>, and Scott D. Bridgham<sup>2</sup>

<sup>1</sup> Chapman University, Orange, CA;

<sup>2</sup> University of Oregon, Eugene, OR;

<sup>3</sup> Florida State University, Tallahassee, FL;

<sup>4</sup> Oak Ridge National Laboratory, Oak Ridge, TN

Contact: [jkeller@chapman.edu](mailto:jkeller@chapman.edu)

BER Program: TES

Project: University Award

Peatlands contain one-third of the world's soil carbon (C). Much of this C is stored deep in the soil profile, where water-logged and anaerobic conditions have allowed C to accumulate for thousands of years. It is currently unknown if these vast C stores will remain belowground or if they will be respired as carbon dioxide (CO<sub>2</sub>) and/or methane (CH<sub>4</sub>) in the face of ongoing global change. Understanding, and modeling, the future fate of this soil C remains a pressing issue in global biogeochemistry. We assessed the response of CO<sub>2</sub> and CH<sub>4</sub> production in a boreal peatland following 13 months of deep-peat warming (DPW), 16 months of subsequent whole- ecosystem warming (surface and deep warming; WEW), and 4 months of elevated atmospheric CO<sub>2</sub> concentrations as part of the Spruce and Peatland Responses Under Changing Environments (SPRUCE) project in northern Minnesota, USA. This project includes 5 temperature treatments that warmed the entire 2 m peat profile from 0 to +9 °C above ambient temperature with and without elevated atmospheric CO<sub>2</sub> concentrations (~850 ppm). Soil cores were collected at multiple depths beneath the peatland surface (-20 to -200 cm) from each experimental enclosure at SPRUCE and anaerobically incubated at in situ temperatures for 1-2 weeks. Following DPW, only CH<sub>4</sub> production from surface depths (e.g., -30 cm) was positively correlated with elevated temperature. However, during WEW, both surface and deep peat CH<sub>4</sub> production increased with rising temperature. There was little indication that elevated CO<sub>2</sub> influenced CH<sub>4</sub> production. Surface peat had greater CH<sub>4</sub> production than deeper peat, implying that increased CH<sub>4</sub> emissions in response to warming observed in the field were largely driven by surface peat dynamics. Radiocarbon analyses suggest that CO<sub>2</sub> and CH<sub>4</sub> are produced from the decomposition of both young and ancient C sources. The CO<sub>2</sub>:CH<sub>4</sub> ratio was inversely correlated with temperature across all depths during WEW, indicating that the entire peat profile is becoming more methanogenic with warming. This result was supported by *in situ* measurements of porewater CO<sub>2</sub> and CH<sub>4</sub> concentrations which displayed the same trend. Thus, our results suggest that the vast C stores at depth in peatlands are responsive to warming only after a significant lag period, but that ecosystem responses remain largely driven by surface peat.